

LORENZ ON LIGHT: A PRECOCIOUS PHOTON PARADIGM

H. C. POTTER

ABSTRACT. In 1867, during the time when Maxwell was publishing his electromagnetic theory, L. Lorenz published his theory equating light vibrations with electric currents. Starting from Kirchhoff's Ohm's law expression, Lorenz introduces scalar potential retardation to obtain an expression for the electric field using retarded potentials. In his theory Lorenz sets the vacuum light speed equal to $\sqrt{2}/2$ times Weber's measured superluminal value for magnetic induction speed. Using the wave equation Green's function paradigm Lorenz reduces his integral, retarded Ohm's law formulation to a differential formulation for current density. This represents his solution to the electromagnetic action at a distance problem. During the reduction he presents a light like, plane wave solution for current density that can not satisfy initial conditions on the expanding wave front and, using a faulty by parts integration that neglects retardation, he develops the eponymic Lorenz condition. Although generally accepted, this condition relates vector and scalar potentials erroneously. A proper relation is given and major consequences are developed in appendices. As a finale Lorenz suggests a light model that I interpret as magnetoinductive waves. Such waves are only now being studied using Kirchhoff's laws without retardation. I redo this important paper using modern day notation with critical annotations. This elevates the paper to a teaching tool position in the millennial search to understand light's dual nature as wave and corpuscle and brings us full circle, back to Kirchhoff's laws where Lorenz began.

Key words and phrases. Light theory, Lorenz condition, Maxwell equations, Photons.

1. INTRODUCTION

In researching photon models I obtained the 1867 article entitled “On the identity of the vibrations of light with electrical currents” [1] expecting to find the rationale for the Lorenz condition that relates electromagnetic potentials. I found much more: wave equations for light derived from Ohm’s law by retarding scalar potential propagation and reference to light as rotationally polarized by transversely circulating currents. I interpret these currents as forming magnetoinductive waves exhibiting autonomous progression. As such, these current loops represent a precocious model for light quanta whose photon name was not proposed [2] for another 59 years. The role Lorenz played in developing electromagnetic theory and, in particular, his 1867 paper precedes gauge theory development. Gauge theory has come to be based on the observation by H. A. Lorentz (1853-1928) that special potential transformations leave select electromagnetic Maxwell fields invariant [3]. This observation was apparently made as an aside when Lorentz discovered that the Lorenz condition was required for the Maxwell electromagnetism equations to be satisfied for fields derived from wave function potentials. The Lorenz condition inverse development from wave function potentials in [1] was found only later and led to shared attribution, but the Lorenz development was not critically scrutinized as is done below. This oversight resulted in a failure to detect that the Lorenz condition development can be deemed flawed when the vector potential is treated as a local function whose spatial derivatives at a point depend only on function values near the point. This failure is etiologically significant; for, if the Lorentz discovery can be inverted, Maxwell’s electromagnetism equations and, ultimately, the Lorentz transformation on which relativity is based would follow from Ohm’s law and Lorenz retardation. This flaw is exposed below by translating the Lorenz equations to modern vector form and explained in 2.5. Its effect on the field equations is presented in Appendix B. In this paper I give a detailed, section by section reprise with annotations appropriate to our modern day understandings. But first some background will help explain why this paper is important and why it was nearly lost to posterity.

Lorenz, Ludvig Valentin, was a Danish physicist who lived from 1829-1891. Since the article was published during the time period (1865-1873) when Maxwell (1831-1879) was publishing his work and does not refer to Maxwell’s formulation, the mathematics needs

translation into present day terms for ready comprehension. With Maxwell's vacuum light velocity prediction $c_0 = 1/\sqrt{\epsilon_0\mu_0}$ from the vacuum electric and magnetic constants for transverse electromagnetic waves, his field equations became the standard for modeling light as traveling plane waves. These Maxwell waves carry in phase electric and magnetic fields orthogonal to one another and the travel direction. The subsequent discovery that light had a particle like character in the early 20th century posed duality reconciliation as a problem for physicists. Attempts to resolve this problem that my search has uncovered are succinctly summarized in Table 1 which supplements and updates the review in [4].

In the next section I present the Lorenz work as I interpret it using present day vector calculus. I present the work in small, well defined sections with most sections followed by my critique. Following this presentation I discuss this work's educational significance more broadly.

2. PRESENTATION

In this section I concisely present the Lorenz analysis using present day symbology. The original equation numbering, sometimes unconventional, is retained to facilitate cross checking. With some clear exceptions many other equations represent unnumbered Lorenz relations. Since the symbol we presently associate with full differentiation is used throughout the work, our symbol for partial differentiation has been introduced where required. Subsection critiques have been added to facilitate concept binding.

2.1. Analytical Basis. Lorenz bases his analysis on a starting formulation that he attributes to Kirchhoff.

2.1.1. Kirchhoff formulation. Consistent with the article title, Lorenz begins with equations that today we would associate with Ohm's law expressed as

$$\mathbf{J} = \kappa\mathbf{E} = -\kappa\left(\nabla\Omega + \frac{2}{c}\frac{\partial\mathbf{A}}{\partial t}\right) \quad (\text{L1})$$

where the current density, $\mathbf{J}$, equals the conductivity, κ , times an electric field, $\mathbf{E}$, equal to a negative scalar field gradient, $-\nabla\Omega$, plus a normalized negative vector potential temporal

variation, $-\frac{2}{c}\frac{\partial \mathbf{A}}{\partial t}$. Following Kirchhoff, Lorenz expressed the vector potential [3] as

$$\mathbf{A}_K = \frac{2}{c} \iiint \frac{\mathbf{R}}{R^3} \mathbf{R} \bullet \mathbf{J}(\mathbf{r}', t) dv' \quad (1)$$

and the scalar potential as Coulomb potential superposition

$$\Omega = \iiint \frac{\epsilon'}{R} dv' + \int \frac{e'}{R} ds' \quad (2)$$

with the continuity equations for charge

$$\left. \begin{aligned} \nabla \bullet \mathbf{J} &= -\frac{1}{2} \frac{\partial \epsilon}{\partial t}, \\ \mathbf{J} \bullet d\mathbf{n} &= -\frac{1}{2} \frac{de}{dt} \end{aligned} \right\} \quad (L2)$$

where the ϵ' is the charge density at the relative position $\mathbf{R} = \mathbf{r} - \mathbf{r}'$ in a Cartesian system and e' is the charge density on the surface element ds' with inward normal $d\mathbf{n}$. Here the primed densities are in electric potential units. So the electric constant with value 8.85E-12 Fm⁻¹, F=QV⁻¹, must be introduced where required.

2.1.2. Kirchhoff Formulation Critique. The Eqs.(L2) express charge conservation. The first follows redundantly from the Ampere law expressed using the Maxwell magnetizing force, $\mathbf{H}$, and electric induction, $\mathbf{D}$, as

$$\nabla \times \mathbf{H} = \frac{\partial \mathbf{D}}{\partial t} + \mathbf{J} \quad (3)$$

with $\nabla \bullet \mathbf{D} = \rho$ since $\nabla \bullet (\nabla \times \mathbf{H}) \equiv 0$. The second follows from $\mathbf{J} = \rho \mathbf{v}$ where ρ is the local charge per unit volume and $\mathbf{v}$ is its local velocity. When the magnetic induction is defined by

$$\mathbf{B} = \frac{2}{c} \nabla \times \mathbf{A}, \quad (4)$$

the Eq.(L1) expresses the Faraday law

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \quad (5)$$

since $\nabla \times \nabla \Omega \equiv 0$. Thus, without explicitly invoking magnetic effects, the formulation is implicitly consistent with Maxwell's four equations. The scalar potential can be considered a Green's function solution for the Poisson equation, $\nabla^2 \Omega = -\rho$, in a connected volume

that includes surface terms that are seldom considered in present day analyses. The vector potential is an *ad hoc* expression. Below it will be brought surprisingly into conformity with the present day retarded form.

The formulation used by Lorenz considers charge and current distributions to be fundamental observables from which real potentials can be obtained. This subverts the less than pure field Maxwell conception that real charge and current distributions give rise to observable electromagnetic fields but nonphysical potentials [5, Chapter XII]. Contributing to potential nonphysicality is susceptibility to transformation with an arbitrary gauge function, χ , that causes

$$\left. \begin{aligned} \mathbf{A} \rightarrow \mathbf{A}' &= \mathbf{A} + \nabla\chi, \\ \Omega \rightarrow \Omega' &= \Omega - \frac{1}{c} \frac{\partial\chi}{\partial t}. \end{aligned} \right\} \quad (6)$$

Many such transformations are presented in [6]. Two are given special recognition. The χ choice that gives $\nabla \bullet \mathbf{A} = 0$ is called the Coulomb gauge, while that that gives $\nabla \bullet \mathbf{A} + \frac{1}{c} \frac{\partial\Omega}{\partial t} = 0$ is called the Lorenz gauge. Expressions for the new potentials have come to be called “gauges” without regard for gauge function existence [7]. This custom is inappropriate, because the gauge expressions can be satisfied by nongauge transformations. For example, the Coulomb gauge relation can be satisfied whenever the vector potential is augmented by some vector’s curl; but, by the Helmholtz Theorem, that augmentation can not be a gauge function’s gradient. Thus, the gauge relations actually restrict the problem relevant potentials. This justifies the proposal in [8] that electromagnetic potential specification is essential for obtaining complete electromagnetic field distributions. This proposal is supported by the finding in 2.5 that there are *two* distinct retarded scalar potentials, Eq.(7) $\overline{\Omega}$ dependent on remote charge and Eq.(21) $\overline{\Omega}_0$ dependent on remote current.

2.2. Basis Extension. Lorenz considers the Eq.(L1) right side terms to be the first terms in series expansions that add potential retardation.

2.2.1. Retardation. Designating the retarded time by $t - \frac{R}{a}$, Lorenz defines the new function

$$\overline{\Omega} = \int \int \int \frac{\epsilon'(t - \frac{R}{a})}{R} dv' + \int \frac{e'(t - \frac{R}{a})}{R} ds'. \quad (7)$$

The charge densities are expanded to give

$$\left. \begin{aligned} \epsilon'(t - \frac{R}{a}) &= \epsilon' - \frac{\partial \epsilon'}{\partial t} \frac{R}{a} + \frac{\partial^2 \epsilon'}{\partial t^2} \left(\frac{R}{a}\right)^2 \frac{1}{2} - \dots, \\ \epsilon'(t - \frac{R}{a}) &= \epsilon' - \frac{d\epsilon'}{dt} \frac{R}{a} + \frac{d^2 \epsilon'}{dt^2} \left(\frac{R}{a}\right)^2 \frac{1}{2} - \dots. \end{aligned} \right\} \quad (8)$$

When the expansions are inserted

$$\nabla \bar{\Omega} = \nabla \Omega + \frac{1}{2a^2} \left[\int \int \int \frac{\mathbf{R}}{R} \frac{\partial^2 \epsilon'}{\partial t^2} dv' + \int \frac{\mathbf{R}}{R} \frac{d^2 \epsilon'}{dt^2} ds' \right] - \dots. \quad (9)$$

Careful coordinate tracking confirms development validity to this point, but note that the integrands no longer depend on the retarded time. Here Lorenz states that using Eqs.(L2) and partial integration gives

$$\nabla \bar{\Omega} + \frac{1}{a^2} \frac{\partial}{\partial t} \int \int \int \frac{\mathbf{J}'}{R} dv' = \nabla \Omega + \frac{c}{2a^2} \frac{\partial \mathbf{A}_K}{\partial t}, \quad (L4)$$

where $\mathbf{J}'$ indicates that $\mathbf{J}$ depends on $t - \frac{R}{a}$ and $\mathbf{r}'$ rather than t and $\mathbf{r}$.

2.2.2. Retardation Critique. Despite the 2.2.1 predominant significance for the Lorenz paper, the $\nabla \bar{\Omega}$ series expansion contains too many terms for the integration by parts hint to facilitate obtaining the final relation. An obscure, present day vector analysis integral theorem provides better direction. The theorem states

$$\int \int \int \mathbf{V}_1 (\nabla \bullet \mathbf{V}_2) dv = \int \mathbf{V}_1 \mathbf{V}_2 \bullet d\mathbf{s} - \int \int \int (\mathbf{V}_2 \bullet \nabla) \mathbf{V}_1 dv. \quad (10)$$

So, with $\mathbf{V}_1 = \frac{\mathbf{R}}{R}$ and $\mathbf{V}_2 = \mathbf{J}$

$$\left. \begin{aligned} \int \int \int \frac{\mathbf{R}}{R} (\nabla \bullet \mathbf{J}) dv &= \int \frac{\mathbf{R}}{R} \mathbf{J} \bullet d\mathbf{s} - \int \int \int (\mathbf{J} \bullet \nabla) \frac{\mathbf{R}}{R} dv, \\ \int \int \int (\mathbf{J} \bullet \nabla) \frac{\mathbf{R}}{R} dv &= \int \int \int \frac{\mathbf{J}}{R} dv - \int \int \int \mathbf{J} \bullet \mathbf{R} \frac{\mathbf{R}}{R^3} dv. \end{aligned} \right\} \quad (11)$$

With these Eq.(L4) is readily verified when signs are adjusted for converting the integration and differentiation variables from $\mathbf{r}$ to $\mathbf{r}'$ in Eq.(11). The Eq.(L4) right hand side is an expansion for which only the first two members have been retained and includes a negligible term required to restore retardation to the current density.

2.3. Retarded Basis.

2.3.1. *Lorenz Formulation.* If a is assumed to equal the vacuum light speed, the neglected Eq.(L4) terms will be negligible for the small distances studied, “a few feet”, provided the second and higher order current component time derivatives be “not very great”. Under these conditions Eq.(L1) can be rewritten using Eq.(L4) to give

$$\mathbf{J} = -\kappa \left(\nabla \bar{\Omega} + \frac{2}{c} \frac{\partial \mathbf{A}_L}{\partial t} \right). \quad (\text{A})$$

Here $\mathbf{A}_L$ has the form

$$\mathbf{A}_L = \frac{2}{c} \int \int \int \frac{\mathbf{J}'}{R} dv'. \quad (12)$$

Using Eq.(L4), Eq.(A) can be written as

$$\mathbf{J} = -\kappa \left(\nabla \Omega + \frac{c}{2a^2} \frac{\partial \mathbf{A}_K}{\partial t} + \left(\frac{4}{c^2} - \frac{1}{a^2} \right) \frac{c}{2} \frac{\partial \mathbf{A}_L}{\partial t} - \dots \right). \quad (13)$$

Setting $2a = c$ recovers Eq.(L1). However, making a “infinitely great” gives an experimentally confirmed form as well. Thus, a must be regarded as a very great undetermined value for which $a\sqrt{2} = c$ would represent a root mean reciprocal square (RMrS) value for these extremes.

2.3.2. *Lorenz Formulation Critique.* For his constant c , Lorenz uses the value

$$c_W = 284736 \text{ miles}/(\text{sec}). \quad (14)$$

Lorenz attributes c_W to experimental determination by Weber for inductive current transfer. He acknowledges that this value is about $\sqrt{2}$ times the then known light speed. For the present day vacuum light speed $c_0 = 299792458 \text{ ms}^{-1}$,

$$\frac{c_W}{c_0} = \frac{284736 \text{ miles}/(\text{sec})}{299792.458 \text{ kms}^{-1}} 1.60935 \text{ km}/\text{mile} = 1.529. \quad (15)$$

The c_W superluminality is explained in [9] as due to a $\sqrt{2}$ scale factor in Weber’s 1851 formulaic definition. Lorenz considers his relation $c_0 = c_W/\sqrt{2}$ as a new light speed determination. He, therefore, takes $a = c/\sqrt{2}$.

Recent measurements [10] confirm superluminal near field magnetic induction propagation speed. There are also several recent measurements for near field longitudinal electric field propagation speed with finite values [11]. These may indicate reality for the v-gauge

[6] that allows arbitrary scalar potential propagation speed if other observable effects could be tied to the gauges. The v-gauge applies to the configuration dependent potentials which determine the measured fields. These measured fields can be gauge independent, but as mentioned in 2.1.2 they need not be. Making the Lorenz condition a scalar function has been proposed as a solution to this impasse [12]. This solution is supported by the 2.5 Eq.(20) and gives field equations identical to the Lorenz equations presented in Appendix B. The Eq.(21) $\overline{\Omega}_0$ gives the scalar function a specific physical definition. To the extent that the v-gauge enters into photonic crystal [13, 14] and microwave lens [15, 16] design, its reality only recently has been subjected to serious study. Formal field propagation speed expressions are developed in Appendix C.

2.4. Wave Representation. Having determined that the retardation speed a is not restrained by the foregoing development, Lorenz proposes to find another method for its determination and for supporting or correcting that development. Although he purports to have found several ways that retardation might be effected, he rejects such hypothecation because its significance would require selecting a most probable hypothesis. This qualm has not been an impediment for subsequent authors, however. For examples see Table 1. Instead, he resolves to further develop the retardation result with the expectation that added insight will follow. To this end, he considers wave propagation for a general source function ϕ using the expression

$$\square_a \int \int \int \frac{\phi(t - \frac{R}{a}, x', y', z')}{R} dx' dy' dz' = -4\pi\phi(t, x, y, z) \quad (\text{L5})$$

where, the d'Alembertian, $\square_a = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} - \frac{1}{a^2} \frac{\partial^2}{\partial t^2}$.

2.4.1. Wave formulation. Applying this to Eq.(A) using the Eqs.(7) and (12) definitions gives

$$\square_a \mathbf{J} = 4\pi\kappa \left(\nabla \epsilon' + \frac{4}{c^2} \frac{\partial \mathbf{J}}{\partial t} \right) \quad (16)$$

with which the first Eq.(L2) relation is associated. Lorenz displays a one transverse current component, two-dimensional solution for $\epsilon = 0$, periodic in one-dimension and attenuated in the other. The published solution is inconsistent with the above relations, because for $\epsilon = 0$ at least two current components are required. However, a suitable one component

solution instance is the following:

$$\left. \begin{aligned} J_x(x, y, z, t) &= e^{-(z)h} \cos[p(\omega t - z)] \\ J_y(x, y, z, t) &= 0 \\ J_z(x, y, z, t) &= 0 \end{aligned} \right\} \quad (\text{L6'})$$

with

$$h^2 a^2 = p^2(a^2 - \omega^2) \quad \text{and} \quad hc^2 = 8\pi\kappa\omega. \quad (\text{L7})$$

Both h and p have inverse length units, while κ has inverse time units obtained from and inverse electric resistivity with F/ms units by electric constant division. As published, Lorenz had an x substituted for z in the exponential.

For Lorenz this preliminary Eq.(A) treatment demonstrates several significant results: periodical electrical currents are possible; these periodical electric currents travel with (phase) velocity, ω ; and they exhibit transverse vibrations, at right angles to the propagation direction. Furthermore, if light vibrations are assumed to be electric currents “ a is the velocity with which electrical action is propagated through space.”

2.4.2. Traveling Wave Critique. The Eq.(L5) form presumes $\phi(0, x, y, z) = 0$ with zero first derivatives as initial conditions [17, pp. 147-148] and applies only in the region $R < at$ and only up to the nearest scattering inhomogeneity.

The Eq.(L6') solution has one critical defect that seems common to sourced wave equation solutions described as traveling waves: It does not, and can not, satisfy the initial conditions anywhere on the solution front at $R = at$. As a consequence, traveling wave solutions to wave equations for one spatial dimension can not be considered valid solutions. But for higher spatial dimensions there may be solutions that incorporate discontinuity at the wave front [17, Ch. IX]. As an alternative ray formulation, Bateman has examined singularity propagation on wave equation characteristics [18]. Being located *on the expanding wave front*, these singularities provide the initial condition transition. I propose that such solutions be called “photon solutions”, because with neither wave packet dispersion [19] nor quantum mechanical non-locality they provide localization that has long eluded discovery [20]. Bateman acknowledges that there is no evidence that his singular fields can be built up

by elementary electromagnetic field superposition. He calls these fields "æthereal", and proposes that their superposition defines the electromagnetic field about moving singularities. [21, Sec. 2]

2.5. Lorenz Condition. For use in transforming his Eq.(A) to his Eq.(B) below, Lorenz develops in the manner next presented what we have come to call the Lorenz condition.

2.5.1. Development. By writing Eqs.(L2) with the retarded time $t - \frac{R}{a}$ substituted for the temporal argument and substituting them into Eq.(7) after t differentiation, he should obtain

$$\frac{\partial \bar{\Omega}}{\partial t} = -2 \iiint \frac{\nabla' \bullet \mathbf{J}'}{R} dv' - 2 \int \frac{\mathbf{J}' \bullet d\mathbf{n}'}{R}. \quad (17)$$

Lorenz then says integration by parts turns the expression into one involving the Eq.(12) vector potential that we call the Lorenz condition

$$\frac{\partial \bar{\Omega}}{\partial t} = -c \nabla \bullet \mathbf{A}_L. \quad (18)$$

2.5.2. Development Critique. I find the transition from Eq.(17) to Eq.(18) troubling. Certainly, the explicit surface integration in Eq.(17) can be suppressed to recover a single volume integration, but factoring out the $\nabla \bullet$ is not obviously valid since inside the integral it acts on the $\mathbf{r}'$ in $\mathbf{J}(\mathbf{r}', t - \frac{R}{a})$ and outside the integral it acts only on $\mathbf{r}$ in R . If, however, the $\nabla \bullet$ acting on $\mathbf{r}$ and $\nabla' \bullet$ acting on $\mathbf{r}'$ are applied to the argument in the Eq.(12) $\mathbf{A}_L$ integral definition and the results added, we obtain

$$-\nabla \bullet \left(\frac{\mathbf{J}'}{R} \right) = \nabla' \bullet \left(\frac{\mathbf{J}'}{R} \right) - \frac{\nabla' \bullet \mathbf{J}'}{R} + \frac{1}{aR^2} \mathbf{R} \bullet \frac{\partial \mathbf{J}'}{\partial t} \quad (19)$$

when $\mathbf{J}' = \mathbf{J}(\mathbf{r}', t - \frac{R}{a})$. Significantly, the temporal derivative from $\nabla \bullet \mathbf{J}'$ remains. It is required to cancel a corresponding term from $\nabla' \bullet \mathbf{J}'$; but that term can not be explicitly displayed without defining a nonphysical, derived residue that neglects retardation, *i.e.* by considering the vector potential to be a non local function whose spatial derivatives are determined by conditions far from the point at which the derivative is to be determined. Now, subjecting Eq.(19) to volume integration gives the Lorenz condition with retardation

$$-c \nabla \bullet \mathbf{A}_L = \frac{\partial \bar{\Omega}}{\partial t} + \frac{\partial \bar{\Omega}_0}{\partial t} \quad (20)$$

when $\int \int \int \nabla' \bullet \left(\frac{\mathbf{J}'}{R} \right) dv' = \int \left(\frac{\mathbf{J}'}{R} \right) \bullet ds'$, by the Gauss theorem, vanishes on the volume boundary, and

$$\bar{\Omega}_0 = \frac{2}{a} \int \int \int \frac{\mathbf{R} \bullet \mathbf{J}'}{R^2} dv'. \quad (21)$$

By applying Eq.(L5) to Eq.(20) $\bar{\Omega}_0$ can be shown to satisfy the homogeneous wave equation, $\square_a \bar{\Omega}_0 = 0$. Thus, $\bar{\Omega}_0$ is a general solution for the configurational scalar potential when fixed charge is absent.

This development is unnecessary for showing Eq.(A) and Eq.(B) equivalent in 2.6.1 below when $\nabla \bullet \mathbf{J} = 0$. However, as Lorenz, himself, confirms in 2.6.1 $\nabla \bullet \mathbf{J} = 0$ is the necessary and sufficient condition for replacing electromagnetic action at a distance, Eq.(A), by strict locality, Eq.(B). This means Eq.(17) is identically zero. So the Eq.(18) Lorenz condition reduces to the Coulomb gauge, $\nabla \bullet \mathbf{A}_L = 0$; but Eq.(20) shows this to be true only when the Eq.(21) temporal derivative also vanishes. So the correction to the Eq.(18) Lorenz condition found here has nontrivial consequences. Some are expounded in the appendices. Importantly, the Eq.(20) Lorenz condition with retardation gives modified field equations developed in Appendix B. Maxwell equation dependence on the Lorenz condition has been examined recently [8, 12].

2.6. Light Equation.

2.6.1. *Wave Reformulation.* Through a complex mathematical manipulation that we would effect by using the vector identity

$$\nabla \times (\nabla \times \mathbf{J}) = \nabla (\nabla \bullet \mathbf{J}) - \nabla^2 \mathbf{J} \quad (22)$$

in Eq.(16) for charge free space, Lorenz succeeds in rewriting his Eq.(A). With his $c = a\sqrt{2}$ the result is his light equation augmented with a conductivity term that introduces attenuation

$$\nabla \times (\nabla \times \mathbf{J}) + \frac{1}{a^2} \frac{\partial^2 \mathbf{J}}{\partial t^2} + 8\pi\kappa \frac{1}{a^2} \frac{\partial \mathbf{J}}{\partial t} = 0. \quad (B)$$

He also shows that the original equations can be recovered through another mathematical manipulation made more complex than the first by the necessity to reintroduce the non-zero charge density gradient. These manipulations are more fully described in Appendix A. During both he explicitly displays the vector potential wave function that follows from the

Eq.(L5) paradigm applied to Eq.(12):

$$\square_a \mathbf{A}_L = -\frac{8\pi}{c} \mathbf{J}. \quad (23)$$

The vibrations described by Eqs.(B) are transverse; longitudinal vibrations will not be possible. The first is confirmed in 2.4.1; the second, because for $\nabla \bullet \mathbf{J} \equiv \Theta$, $\nabla \bullet$ Eq.(B) gives

$$\frac{\partial \Theta}{\partial t} + 8\pi\kappa\Theta = 0 \quad (24)$$

after one time integration. Clearly Θ can not be periodic and $\lim_{t \rightarrow \infty} \Theta = 0$. Since Eqs.(L2) give $\Theta = -\frac{1}{2} \frac{\partial \epsilon}{\partial t}$, no free charge “development” is possible within a conductor.

2.6.2. Wave Reformulation Critique. As discussed more fully in Appendix B, Lorenz claims to have solved the self field, action at a distance problem inherent in Eq.(A) with $\mathbf{A}_L$ defined by Eq.(12). The Eq.(B) which contains only one, local field is his solution. The Appendix B also shows that Eq.(B) is not required to obtain the field equations from the foregoing Lorenz development when Eq.(4) is used to define the magnetic induction. Lorenz, himself, never gives an expression for the magnetic field. He only mentions that its propagation speed is apparently superluminal as covered in 2.3.2. Electromagnetic field retardation is discussed more fully in Appendix C.

2.7. Concluding Remarks. Lorenz ends the work with paragraphs giving a summary, a suggestion for extending the development to heterogeneous material, an æther dismissal and, of course, his alternative conception for light vibration.

2.7.1. Lorenz Conclusion. Lorenz considers the work to be a “new proof” that light vibrations are electrical currents. He claims to have shown that vibrations described by Eq.(B), his “laws of light” augmented by absorption, are electric currents described by Eq.(A). This result is said to depend on no assumed physical hypothesis.

Lorenz deems extension to heterogeneous bodies best done by starting from the differential equations, possibly an indirect reference to the Maxwell relations given in 2.1.2, while regarding a and κ to be variable quantities. Lorenz laments that the Eq.(A) simplicity could not be preserved. So Eq.(A) must be considered as strictly applying to homogeneous bodies. The important result, however, is that “electrical forces require time to travel.” As a result,

the action at a distance, non-locality, contained in Eq.(A) in fact results from local action described by Eq.(B).

Since the æther theory for light at the time was one with æther particles in motion, the Lorenz conception that light vibrations were electric currents implied æther particles to be electric in nature. So electric currents would be described by progressive æther particle motion. Lorenz found this prospect untenable. So, rather than reject his electric current association with light vibrations, he found it expedient to reject the æther concept.

The linearly transverse current solution given by Eq.(L6') gives Lorenz no reason to prefer his electric current model for light vibrations over the æther model. But he justifies the preference by proposing that the currents are actually rotary. In good conductors current circulates in the rotation direction, but in bad conductors it is propagated by induction in a direction perpendicular to the planes containing the periodical currents.

2.7.2. Conclusion Critique. For Lorenz, the conception that electrical interactions require time for propagation is so obvious and scientifically acceptable that it may be assumed as fact. With this assumed physical hypothesis, Lorenz has bypassed the Maxwell field formulation that we consider fundamental.

Although Lorenz states that he had previously “adverted” to the rotary light coupling, he presents no explicit example in his paper. However, the proposal can be looked upon as magnetoinductive coupling. This effect has been demonstrated recently to transfer 60 watts with 40% efficiency between self-resonant coils with resonant frequency near 10 MHz over a distance exceeding 2 meters in a strongly coupled regime [22]. This power transfer does not have simple Poynting character. If the energy were to propagate by inductive regeneration as proposed by Lorenz, the rotary electric field would progress autonomously with alternating electric field direction. This possibility might be experimentally evaluated in the wireless power transfer apparatus by attempting to measure the induced electromotive force phase and power transfer efficiency change produced by adding one or more resonant coils between the source and receiver coils. These waves add to self-trapping [23] and synchrotron electric fields [24, Figure 5] as possible Maxwell wave alternatives.

For photons Dirac [5] has derived a phase space volume equal to the Planck constant cubed. With momentum equipartition this gives an effective spatial volume per photon

equal to $[3\lambda]^3$, each containing the photon energy $h\nu$. Magnetosphere continuous pulsations at 0.005 Hz investigated by satellite [25] have revealed antiphase electric and magnetic fields within the theoretical spatial volume equal to $1.1 \times 10^{24} \text{ km}^3$, about 10^6 sun volumes. With a linear dimension equal to 10^8 km , passage at vacuum light speed in 5 minutes or less agrees well with the reported pulse durations. If these observations can be attributed to individual photons, a new study regime may be opened.

3. DISCUSSION

Frequently in attempting to solve complex problems, we find ourselves retracing paths that fail to provide an adequate solution. Table 1 shows that our search for a physical photon model is an example. In such cases, a fresh start can often help to break free from the limitations that impede progress. The fresh start can generally be either closely examining prior attempts for possible flaws or taking an entirely different approach. In subjecting the 1867 Lorenz paper to detailed, critical examination, this paper applies both these remedies for failed attempts to resolve light wave/particle duality.

Table 1 examination shows, and [20] confirms, that nearly all attempts to describe particle like light behavior are based on the Maxwell electromagnetism formulation. Being written before Maxwell attained unquestioned authority, the 1867 Lorenz paper begins at the well established Ohm's law level as formulated by Kirchhoff, Eq.(L1), with the electric field defined by two potentials, a scalar potential and a vector potential. By introducing retardation, apparently a novel idea at the time, to eliminate potential tracking instantaneously with source position [26], Lorenz obtains his Eq.(A) Ohm's law formulation. Using Eq.(13) Lorenz demonstrates that the retardation speed is not theoretically established. By choosing $a = c_0 = c_W/\sqrt{2}$, however, he interrelates the propagation speeds for inductive current transfer and for light. This choice for scalar potential propagation speed was experimentally validated only recently [11, Tsontchev].

To obtain more insight into electric signal propagation, Lorenz applies the Green's function solution for the sourced wave equation, Eq.(L5), to his retarded potential Ohm's law formulation, Eq.(A). The result, Eq.(16), is effectively a sourced wave equation for the

electric field. Lorenz presents a solution for this equation in charge free space. After typographical error correction, the solution represents a plane polarized, traveling wave exhibiting light like properties by being periodic and transverse to the propagation direction. In 2.4.2 I note that traveling waves do not satisfy initial conditions on the expanding solution boundary. But there appear to be singular wave equation solutions that can satisfy the initial conditions on the expanding wave front [18]. Such solutions may represent the elusive localized photon.

For use in transforming Eq.(16) into a form that he had previously published as an equation for light, Lorenz develops what we call the Lorenz condition, Eq.(18), from Eq.(17). Flaws are exposed in 2.5 and 2.6. The 2.5 development must be corrected to avoid inadvertent retardation suppression in the vector potential current density argument. This correction yields field equations derived in Appendix B with an augmented Maxwell form. Furthermore, in the 2.6 development, he then shows that $\nabla \bullet \mathbf{J} = 0$. So, from Eq.(17) the scalar potential temporal time derivative vanishes and the Lorenz condition without correction, Eq.(18), reduces to $\nabla \bullet \mathbf{A}_L = 0$, the Coulomb gauge.

Only recently has the magnetoinductive wave light model proposed by Lorenz been subjected to observational [27] and theoretical analysis [28]. These models are formed from inductively coupled, capacitively loaded wire rings arranged in 1, 2 and 3 dimensional arrays to simulate common optical phenomena as multiport networks. These elements can not, however, be described by the Lorenz Eq.(B), because the capacitive loading represents free charge development. We do not yet know if the Appendix B Lorenz fields will support magnetoinductive waves, but the requisite structure may be plasma emergent [29, VI.]. More conventionally, however, the capacitor electric field could be taken to describe a wave polarization orientation that varies with capacitor azimuth. Considered as a series connected LC loop, each module would exhibit a representative capacitor charge, $Q_C = CV_0 \cos \omega t$, where $\omega^2 = (LC)^{-1}$ for initial capacitor voltage V_0 and loop inductance L . The loop current, $i_C = -\omega V_0 \sin \omega t$, would produce a time dependent magnetic field through the loop coupling it to its near neighbors with some delay. The electric field defined by the capacitor voltage, $V_C = Q_C/C = V_0 \cos \omega t$, clearly differs in phase from the developed magnetic field, proportional to i_C , by $\pm \frac{\pi}{2}$. To describe general electromagnetic phenomena, these modules would have to be vectorized to properly represent their azimuthal and axial orientation.

With this and the axial magnetic field, the modules can be linearly configured to display wave modes presented in Table 3. The Kirchhoff network waves displayed in Table 3 exhibit modes that are generally excluded from consideration for Maxwell waves and represent a proper complement to the Lorenz legacy.

4. CONCLUSION

The Lorenz 1867 paper has both blemishes and exceptional innovations. The most obvious and disconcerting blemish is the typographical error in Eq.(L6'). The erroneous solution to his Eqs.(16) casts doubt on the whole paper even though the minor revision provided in Eq.(L6') does work. By modern standards his equating light vibrations to electrical currents rather than electric fields is archaic, but future magnetoinductive wave studies may well prove his usage prescient. His Eq.(B) development from Eq.(A) using the effectively null relation that we associate with the Lorenz condition, Eq.(18), rather than proceeding directly from his Eqs.(16) is especially egregious. Most likely he did not want to explicitly set the fixed charge in Eqs.(16) to zero, but found he could do it implicitly with Eq.(18). The 2.5.2 Eq.(20) retarded Lorenz condition reveals a heretofore unrecognized scalar potential, $\overline{\Omega}_0$, developed by currents, Eq.(21). The most pernicious problem, one that persists to this day, is taking the traveling wave solution to the wave equation as a complete, valid solution for wave propagation. This solution by itself does not satisfy the initial conditions on the expanding wave front and is not, therefore, a complete, valid solution. In his book [17], Bateman gives this problem serious consideration, but his work is unfinished and largely ignored as incoherent [30]. A major contribution to our science could result from continuing his work.

The most innovative Lorenz contribution is effective Maxwell equation development, see 2.1.2 and Appendix B, from Kirchhoff's expression for Ohm's law by introducing scalar potential retardation. Although not an astounding innovation, the retained source terms and real value functions with retarded time arguments are exemplary. Now days we commonly suppress sources and use traveling wave phase without considering how the far field waves are to emerge from the retarded near fields. Another modern day time saver uses the complex exponential for periodic functions. By reducing time dependence to a multiplicative factor, this expedient conceals the emergence problem by a subtle change in spatial argument from

$\frac{R}{a}$ to $\frac{\mathbf{k} \bullet \mathbf{R}}{\omega}$ and blatantly confounds Helmholtz Theorem components with longitudinal and transverse fields [20, Sec. 3]. The magnetoinductive wave discussion with reference to Table 3 shows these expedients can mask real phase and orientation relationships. In starting from Ohm's law, Lorenz has implicitly overcome the long standing myopic focus on "visible light." Extending over a 10^{30} factor for frequency and wavelength, Table 2 displays only a minute range for exploration. The lowest, [25], and highest, [35], are science frontiers. The first may indicate that studying photons from the inside may actually be possible as indicated in 2.7.2.

Lorenz most prescient contribution may be his paper ending suggestion that light is rotationally polarized, *i.e.* the electric field is azimuthally directed in cylindrical coordinates rather than radially directed, and propagates by induction along the cylinder axis. This suggestion I have interpreted to mean light actually has a magnetoinductive wave character. Lorenz could not properly develop this concept from his most favored Eq.(B) light wave representation, because magnetoinduction appears to require the free charge in Eqs.(16) and, thus, Eq.(A). Whether such waves are possible with the Appendix B Lorenz fields and Bateman localization is an important subject for future study. But, now, after more than a century, there are signs [28] that magnetoinduction may provide a tool for advancing our long standing bond to Maxwell.

The $\bar{\Omega}_0$ term in 2.5.2 comes from a by parts integration in which the vector potential is treated as a local function with spatial derivatives at a point determined by proximate function values. In reviewing this paper I found that the Eq.(18) Lorenz condition can be developed from the non local vector potential. This follows from applying $\nabla \bullet$ to Eq.(23) and using the Eq.(L2) charge continuity condition to obtain

$$\square_a \nabla \bullet \mathbf{A}_L = -\frac{8\pi}{c} \nabla \bullet \mathbf{J} = \frac{4\pi}{c} \frac{\partial \epsilon}{\partial t}. \quad (25)$$

So applying Eq.(L5) to Eq.(7) gives Eq.(18) to within an additive homogeneous wave equation solution. Thus, the Lorenz condition to be a charge continuity condition restatement for wave function potentials. It allows the field equations to be instructively developed when the potentials treated consistently [31]. In this development, the homogeneous wave function gives nongauge polarization fields. For potentials with independent plane wave vector components, these can emerge from retarded Maxwell fields as light like rays that propagate

autonomously by conserving charge. Alternative treatments using separated-potentials are listed in [32].

5. ACKNOWLEDGMENTS

I would like to thank the Louisville Free Public Library interlibrary loan staff for their valiant efforts in obtaining many references that have made this work possible and especially Teresa Frost whose exceptional facility management helped to provide a felicitous work environment. I would also like to thank an anonymous American Journal of Physics reviewer for suggesting [30].

APPENDIX A. EQ.(A) AND EQ.(B) EQUIVALENCE

As mentioned in 2.6.1, Lorenz presents arguments to show that his Eq.(A) and Eq.(B) are equivalent. The argument from Eq.(A) to Eq.(B) skirts the intermediate Eq.(16). If, however, the vector identity Eq.(22) is applied to Eq.(16), Eq.(B) is equivalent to Eq.(16) and, therefore, to Eq.(A) if and only if $\nabla(\nabla \bullet \mathbf{J}) = 4\pi\kappa \nabla \epsilon'$. Thus, using Eq.(L2), equivalence demands

$$\frac{\partial \nabla \epsilon'}{\partial t} + 8\pi\kappa \nabla \epsilon' = 0. \quad (26)$$

This complements the related Eq.(24) for $\nabla \bullet \mathbf{J}$. Apparently, Lorenz has demonstrated Eq.(26) validity without ever explicitly stating it. To see how, I here recapitulate the Lorenz argument for obtaining Eq.(B) from Eq.(A).

From Eq.(A)

$$\frac{-1}{\kappa} \frac{\partial \mathbf{J}}{\partial t} = \nabla \frac{\partial \bar{\Omega}}{\partial t} + \frac{2}{c} \frac{\partial^2 \mathbf{A}_L}{\partial t^2}. \quad (27)$$

Using Eq.(23) with $a\sqrt{2} = c$, this becomes

$$\frac{1}{2\kappa} \frac{\partial \mathbf{J}}{\partial t} + 4\pi\mathbf{J} = \frac{-1}{2} \left(\nabla \frac{\partial \bar{\Omega}}{\partial t} + c \nabla^2 \mathbf{A}_L \right). \quad (28)$$

Applying the vector identity Eq.(22) gives

$$\frac{1}{2\kappa} \frac{\partial \mathbf{J}}{\partial t} + 4\pi\mathbf{J} = \frac{a}{\sqrt{2}} \nabla \times (\nabla \times \mathbf{A}_L) - \frac{1}{2} \nabla \left(\frac{\partial \bar{\Omega}}{\partial t} + c \nabla \bullet \mathbf{A}_L \right). \quad (\text{L8})$$

Going back to Eq.(A), Lorenz then notes that

$$\nabla \times \mathbf{J} = \frac{-\sqrt{2}\kappa}{a} \frac{\partial \nabla \times \mathbf{A}_L}{\partial t}. \quad (\text{L9})$$

Combining $\frac{\partial}{\partial t}$ Eq.(L8) with $\nabla \times$ Eq.(L9) to eliminate $\mathbf{A}_L$ gives Eq.(B) when the Eq.(18) Lorenz condition is used. But clearly, Eq.(L8) retains a $\frac{\partial \bar{\Omega}}{\partial t}$ term when the Eq.(20) Lorenz condition with retardation is used. By Eq.(24), forcing this residual term to zero effectively forces the free charge development to zero. As this results from using the Eq.(18) in Eq.(L8), the Lorenz condition must be erroneous when the vector potential is considered to be a function with spatial derivatives determined by function values in a small region about the point where the derivative is being determined. The reason is retardation neglect in the by parts integration for its development as discussed in 2.5.

To show their equivalence, Lorenz also demonstrates that Eq.(A) can be obtained from Eq.(B) when Eq.(B) is taken to describe the retarded potential field about some finite volume within which the current actually resides, *i.e.* in Eq.(B) $\mathbf{A}_L$ can replace $\mathbf{J}$! Lorenz justifies this substitution term wise by repeated by parts integration over the conduction volume with zero interface current. When Eq.(B) has been rewritten with $\mathbf{A}_L$ substituted for $\mathbf{J}$, Eqs.(23) and (18) can be used to obtain the form

$$\frac{1}{2} \nabla \frac{\partial \bar{\Omega}}{\partial t} = 4\pi \mathbf{J} + \frac{4\sqrt{2}\pi\kappa}{a} \mathbf{A}_L. \quad (29)$$

Eliminating $\bar{\Omega}$ from Eq.(29) by pairs yields Eq.(L9) for $\nabla \times \mathbf{J}$. Now, substituting Eq.(L9) into Eq.(B) and integrating temporally gives Eq.(L8) from which Eq.(27) can be obtained by reversing the process from which Eq.(L8) was obtained from Eq.(27). One last temporal integration then recovers Eq.(A).

Biologically, the above Lorenz developments may be looked upon as hermaphroditic or even incestuous, because they exhibit unusual recursion. This characteristic alone makes their validity suspect. This suspicion is confirmed in Appendix B. Fortunately, although expressing less physical configuration dependence the modern day vector analysis formalism does validate the final result with the specific restriction that $\nabla \bullet \mathbf{J} = 0$. So in addition to showing how mathematical advances can enhance insight, this example may provide a case study for evaluating proof theory.

APPENDIX B. FIELD EQUATIONS

As stated in 2.1.2 the Lorenz development appears to be consistent with the Maxwell equations when these equations are taken to be strictly local. On this basis, then, the

Maxwell equations are simply descriptive, not predictive. To be predictive the equations must, per Lorenz, reflect retardation. In his 1867 paper, Lorenz considers scalar potential retardation as expressed by Eq.(7). From this he obtains the retarded, local Eq.(L1) electric field as presented in 2.2.1 Eq.(L4) for $2a = c$. His Eq.(A) predicts the local current field determined by the retarded scalar and vector potentials with the latter given by Eq.(12). Lorenz never mentions the magnetic field. Per 2.1.2, the Faraday law for local fields would require the local magnetic field be given by Eq.(4).

Lorenz considers the integral definitions for the retarded potentials to be Green's function solutions for Eq.(L5) wave equations. Properly, these have vanishing values and first temporal derivatives on the expanding solution front. Lorenz solves the self-field or action at a distance problem by eliminating all but one field from his mathematical expressions. In particular, he rewrites his Eq.(A), which interrelates $\mathbf{J}$, $\bar{\Omega}$ and $\mathbf{A}_L$, as his Eq.(B), which involves only $\mathbf{J}$. If this were valid, then Eq.(A) could be written using only $\bar{\Omega}$ or $\mathbf{A}_L$ as well. This can be effected for $\mathbf{A}_L$ by using the Eq.(18) Lorenz condition and Eq.(23), but by the Eq.(L5) Green's function paradigm $\mathbf{A}_L$ is nonlocal. This confirms the suspicion expressed in Appendix A that replacing $\mathbf{J}$ in Eq.(B) with $\mathbf{A}_L$ might not be valid.

Since the single function differential relations for $\bar{\Omega}$ and $\mathbf{A}_L$ are intractable, their integral definitions, Eqs.(7) and (12), along with the augmented Lorenz condition, Eq.(20), and Eq.(A) defining $\mathbf{J}$ and $\mathbf{E}$ and Eq.(4) defining $\mathbf{B}$ define the Lorenz electromagnetic theory. The retarded Maxwell equations can be derived from these:

$$\nabla \bullet \mathbf{B} = \frac{2}{c} \nabla \bullet (\nabla \times \mathbf{A}_L) = 0; \quad (\text{M1})$$

$$\nabla \bullet \mathbf{E} = -\nabla \bullet \left(\nabla \bar{\Omega} + \frac{2}{c} \frac{\partial \mathbf{A}_L}{\partial t} \right) = 4\pi\epsilon' + \left(\frac{2}{c^2} - \frac{1}{a^2} \right) \frac{\partial^2 \bar{\Omega}}{\partial t^2} + \frac{2}{c^2} \frac{\partial^2 \bar{\Omega}_0}{\partial t^2}; \quad (\text{M2})$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}; \quad (\text{M3})$$

$$\begin{aligned} \nabla \times \mathbf{B} &= \frac{2}{c} \nabla \times (\nabla \times \mathbf{A}_L) = \\ &= \frac{2}{c^2} \left[\frac{\partial \mathbf{E}}{\partial t} + 8\pi\mathbf{J} + c \left(\frac{2}{c^2} - \frac{1}{a^2} \right) \frac{\partial^2 \mathbf{A}_L}{\partial t^2} - \frac{\partial \nabla \bar{\Omega}_0}{\partial t} \right]. \end{aligned} \quad (\text{M4})$$

The Eq.(M4) development first uses Eq.(22) and then Eqs.(20) and (23) before finally using the Eq.(A). These simplify considerably with the $c = a\sqrt{2}$ Lorenz assignment. But even

then Eqs.(M2) and (M4) retain a residual, nonlocal Eq.(21) $\overline{\Omega}_0$ dependence. Like the field equations for moving charges that supplement the conventional Maxwell displacement current with a dielectric convection current [33], the Eqs. (M1)-(M4) have the displacement current supplemented by a term due to $\overline{\Omega}_0$ in Eq.(M4). But $\overline{\Omega}_0$ also modifies the charge density in Eq.(M2).

APPENDIX C. ELECTROMAGNETIC FIELD RETARDATION

In his 1867 paper, Lorenz introduced scalar potential retardation to obtain his Eq.(A) for the current density $\mathbf{J}$, effectively the electric field $\mathbf{E}$. Eq.(A) has the same form as Eq.(L1), but now depends on local, retarded scalar and vector potentials given by Eqs.(7) and (12). Since $\mathbf{A}_L$ is derived from $\mathbf{J}$, however, Eq.(A) has the self-field, action at a distance, problem. Lorenz resolves this problem with his Eq.(B) that depends only on $\mathbf{J}$. So Eq.(B) can be taken to describe electric field propagation. Eq.(B) is not a wave equation and comes with the condition that $\nabla \bullet \mathbf{J} = 0$, or $\nabla \bullet \mathbf{E} = 0$ when $\kappa \neq 0$. For consistency, then, the $\nabla \bullet \mathbf{E} = 0$ condition must be retained when $\kappa = 0$. Thus, the electric field propagation speed will be a and the $\nabla \bullet \mathbf{E} = 0$ condition will assure there is no electric field in the propagation direction, no longitudinal electric field.

The Lorenz field equations in Appendix B do not have the $\nabla \bullet \mathbf{J} = 0$ restriction. From $\nabla \times$ Eq.(M3) and Eq.(M4) the Eq.(B) equivalent is found to be

$$\nabla \times (\nabla \times \mathbf{E}) + \frac{2}{c^2} \frac{\partial^2 (\mathbf{E} - \nabla \overline{\Omega}_0)}{\partial t^2} + \frac{16\pi}{c^2} \frac{\partial \mathbf{J}}{\partial t} = 0 \quad (30)$$

when the higher order temporal derivatives are neglected as possibly having a zero coefficient. If $\mathbf{E} - \nabla \overline{\Omega}_0$ in Eq.(30) were mistakenly taken as $\mathbf{E}$, formally the effective propagation speed would satisfy the relation

$$C_{\mathbf{E}}^2 \left| \frac{\partial^2 (\mathbf{E} - \nabla \overline{\Omega}_0)}{\partial t^2} \right| = c^2 \left| \frac{\partial^2 \mathbf{E}}{\partial t^2} \right|. \quad (31)$$

In principle $C_{\mathbf{E}}^2$ would be direction and location dependent and could be greater than c^2 . This argument for a possible superluminal propagation speed cannot be extended to $\nabla \times (\nabla \times \mathbf{B})$, because the $\nabla \overline{\Omega}_0$ in Eq.(M4) drops out. But the argument does apply to $\nabla \times \mathbf{B}$, itself, to

give

$$C_{\nabla \times \mathbf{B}}^2 \left| \frac{\partial(\mathbf{E} - \nabla \bar{\Omega}_0)}{\partial t} \right| = c^2 \left| \frac{\partial \mathbf{E}}{\partial t} \right|. \quad (32)$$

REFERENCES

- [1] L. Lorenz, "On the Identity of the Vibrations of Light with Electrical Currents," The London, Edinburgh and Dublin Philosophical Magazine and Journal of Science Series 4 **34**(1867)287-301.
- [2] G. N. Lewis, "The Conservation of Photons," Nature **118**(1926)874-875.
- [3] J. D. Jackson and L. B. Okun, "Historical roots of gauge invariance," Rev. Mod. Phys. **73**(2001)663-680.
- [4] R. Kidd, J. Ardin and A. Anton, "Evolution of the modern photon," Am. J. Phys. **57** : 1(1989)27-35.
- [5] P A M Dirac , The principles of quantum mechanics, 4th ed., (rev.), Clarendon Press, Oxford University Press (United Kingdom), 1981.
- [6] J. D. Jackson, "From Lorenz to Coulomb and other explicit gauge transformations," Am. J. Phys. **70**:9(2002)917-928 and <http://arxiv.org/abs/physics/0204034>.
- [7] J. A. Heras, "How the potentials in different gauges yield the same retarded electric and magnetic fields," Am. J. Phys. **75**:2(2007)176-183.
- [8] V. V. Onoichin, "On non-equivalence of Lorentz and Coulomb gauges within classical electrodynamics," <http://arxiv.org/abs/physics/0111017>.
- [9] K. S. Mendelson, "The story of c," Am. J. Phys. **74**:11(2006)995-997.
- [10] A. L. Kholmetskii *et al.*, "Experimental test on the applicability of the standard retardation condition for bound magnetic fields," J. Appl. Phys. **101** : 2(2007)3532-1/11.
- [11] C. Monstein and J. P. Wesley, "Observation of scalar longitudinal electrodynamic waves," Europhys.Lett. **59** : 4(2002)514-20; R. I. Tsontchev, A. E. Chubykalo and J. M. Rivera-Juarez, "Coulomb Interaction Does Not Spread Instantaneously," Hadronic Journal **23**(2000)401-24; G.F. Ignatiev and V.A. Leus, "On a superluminal transmission at the phase velocities," in *Instantaneous Action at a Distance in Modern Physics: "Pro" and "Contra"*, edited by A. E. Chubykalo, V. Pope and R. Smirnov-Rueda, Nova Science Pub Inc, 1999, p.203-7.
- [12] K. J. van Vlaenderen, "A generalisation of classical electrodynamics for the prediction of scalar field effects," arXiv:physics/0305098v1 [physics.class-ph].
- [13] E. Ozbay, "Plasmonics: Merging Photonics and Electronics at Nanoscale Dimensions," Science **311**(2006)189-193.
- [14] H. J. Lezec, J. A. Dionne and H. A. Atwater, "Negative Refraction at Visible Frequencies," Science **316**(2007)430-2.
- [15] R. A. Shelby, D. R. Smith and S. Schultz, "Experimental Verification of a Negative Index of Refraction," Science **292**(2001) 77-9.
- [16] T. J. Yen *et al.*, "Terahertz Magnetic Response from Artificial Materials," Science **303**(2004) 1494-6.

- [17] H. Bateman, The mathematical analysis of electrical and optical wave-motion on the basis of Maxwell's equations, Cambridge University Press, 1915.
<http://www.archive.org/details/mathematicalanal00baterich>
- [18] H. C. Potter, "Bateman On Light: Synchrotron Rays," arXiv:0803.3871 [physics.gen-ph].
- [19] C. G. Darwin, "The Uncertainty Principle," Science **73**(1931)653-660.
- [20] O. Keller, "On the theory of spatial localization of photons," Physics Reports **411**:1-3(2005)1-232.
- [21] H. Bateman, "The structure of the æther," Bull. Amer. Math. Soc. **21**:6(1915)299-309.
- [22] A. Kurs *et al.*, "Wireless Power Transfer via Strongly Coupled Magnetic Resonances," Science **317**(2007)83-86.
- [23] A. H. Piekara, "Self-trapping light waves and quanta: Towards an explanation of wave-photon duality," Optics & Laser Technology **14**:4(1982)207-212 .
- [24] J. H. Hannay and M. R. Jeffrey, "The electric field of synchrotron radiation," Proc. Roy. Soc. A **461**(2005)3599-3610.
- [25] P. J. Chi and C. T. Russell, "Phase skipping and Poynting flux of continuous pulsations," J. Geophys. Res. **103**(1998)29479-29491.
- [26] R. Smirnov-Rueda, "Were Hertz's 'crucial' experiments on propagation of electromagnetic interaction conclusive?" Instantaneous Action at a Distance in Modern Physics: "Pro" and "Contra", A. E. Chubykalo, V. Pope and R. Smirnov-Rueda, Eds., Nova Science Pubs., Inc., Commack, NY, 1999.
- [27] M. C. K. Wiltshire *et al.*, "Dispersion characteristics of magneto-inductive waves: comparison between theory and experiment," Electronics Letters **39**:2(2003)215-217.
- [28] R. R. A. Syms *et al.*, "Magneto-inductive waveguide devices," IEE Proc. Micr. Ant. Propag. **153**(2006)111-121.
- [29] R. M. Kinney *et al.*, "Filamentary Magnetohydrodynamic Plasmas," Phys. Plasmas **1**:2(1994)260-280.
- [30] A. Erdelyi, "Harry Bateman. 1882-1946," Obituary Notices of Fellows of the Royal Society, **5**:15(1947)590-618
- [31] H. C. Potter, "Electromagnetic Lorenz Fields," arXiv:0810.1172 [physics.gen-ph].
- [32] O. Keller, "Near-field photon wave mechanics in the Lorenz gauge," Phys. Rev. **76**:6(2007)062110-1/11.;
A. E. Chubykalo and R. Smirnov-Rueda, "Action at a distance as a full-value solution of Maxwell equations: The basis and application of the separated-potential method," Phys. Rev. E **53** : 1(1996)5373-81.
- [33] A. E. Chubykalo and R. Smirnov-Rueda, "Convection Displacement Current and Generalized Form of Maxwell-Lorentz Equations," Mod. Phys. Lett. A **12** : 1(1997)1-24.
- [34] G. N. Lewis, "The Nature of Light," PNAS **12**(1926)22-29.
- [35] A. De Roeck, "Physics of a $\gamma\gamma$, $e\gamma$ and e^-e^- Option for a Linear Collider," arXiv:hep-ph/0311138v1.

TABLE 1. Photon Models

The table entries are keys to appropriate references.

Author	Date	Description
Fermat	1601-65	Space-time geodesic
J. C. Maxwell	1865+	Field equations
L. Lorenz [1]	1867	Inductive, autonomous progression
H. Bateman [17]	1915±	Singular wave functions
A. Einstein	1916	Nadel Strahlung
H. Bateman	1921	Electric doublet
G. N. Lewis [34]	1926	Relativistic proper-time reality
P. A. M. Dirac [5]	1958	Second Quantization and Quantum Electrodynamics
W. H. Bostick	1961	Coiled current loop
T. W. Marshall	1963	Random Electrodynamics
R. K. Nesbet	1971	Semiclassical radiation theory
J. des Cloizeaux	1973	Collective Fermi field excitation
A. M. Arthurs	1979	On electromagnetic field quantization
R. Woodruff & C. Graden	1980	Conceptual zwitter oscillations
T. H. Boyer	1980	Stochastic Electrodynamics
A. H. Piekara[23]	1982	Electromagnetic field self-trapping
M. W. Evans <i>et al.</i>	1995±	Longitudinal magnetic field
J. E. Sipe	1995	Photon wave functions
H. Marmanis	1998	Navier-Stokes equations
D. H. Kobe	1999	Relativistic second quantization
I. V. Lindell	2000	Radiation operator
P. Ghose <i>et al.</i>	2001	Relativistic quantum mechanics
B I & V B Makshantsev	2001	Vector Potential
V. P. Dmitriyev	2003	Hydrodynamic vortex
Hannay & Jeffrey [24]	2005	Synchrotron electric field
P. N. Kaloyerou	2005±	Hidden-variable quantum mechanics
I. Bialynicki-Birula	2006±	Quantum particle
D. Dragoman	2006	Complex electromagnetic fields
D. F. Roscoe	2006	Generalized Maxwell equations
R. R. A. Syms [28]	2006	Magneto-inductive waveguide
B. J. Smith & M. G. Raymer	2007	Photon wave functions

TABLE 2. Photon Characteristics

*Cosmic Microwave Background at 3K equals 0.26 meV.

Energy $h\nu$	Frequency ν Hz	Wavelength λ cm	Intensity $(\frac{h\nu}{\lambda^2})[5]$ $W/(m^2\Delta\nu)$
1Tev	0.241E27	12.4E-17	1.0E29
1Gev [35]	0.241E24	12.4E-14	1.0E20
1Mev	0.241E21	12.4E-11	1.0E11
1Kev	0.241E18	12.4E-8	1.0E2
1ev	0.241E15	12.4E-5	1.0E-7
1mev*	0.241E12	12.4E-2	1.0E-16
1 μ ev	0.241E9	12.4E1	1.0E-25
1nev	0.241E6	12.4E4	1.0E-34
1pev	0.241E3	12.4E7	1.0E-43
1fev	0.241	12.4E10	1.0E-52
1aev [25]	0.241E-3	12.4E13	1.0E-61

TABLE 3. Magnetoinductive Wave Modes

For the first three modes, the loops lie in the paper plane. For the last mode, the loops are normal to a common axis in the paper plane. The magnetic field is considered normal to the loop plane. The electric field is considered to lie in the loop gap. Mode designations are: T, transverse; L, longitudinal; M, magnetic; E, electric.

Mode	Phase	Loop Orientation
TME	$\pm \frac{\pi}{2}$	... C C C C C C ...
TMTLE	$\pm \frac{\pi}{2}$	... J J J J J J ...
TMLE	$\pm \frac{\pi}{2}$	... U U U U U U ...
TELM	$\pm \frac{\pi}{2}$	